

II. Examination Statistics from feather mites living in feather quills

Examination period 15.12.1999 - 31.12.2009

1	Bar-tailed Godwit (<i>Limosa lapponica</i>)	2
2	Beavan`s Bullfinch (<i>Pyrrhula erythaca</i>)	2
3	Black-headed Gull (<i>Larus ridibundus</i>)	1
4	Black-tailed Godwitt (<i>Limosa limosa</i>)	2
5	Blue-naped Mousebird (<i>Urocolius macrourus</i>)	1
6	Chaffinch (<i>Fringilla coelebs</i>)	1
7	Common Redshank (<i>Tringa totanus</i>)	1
8	Eurasian Bullfinch (<i>Pyrrhula pyrrhula</i>)	7
9	European Robin (<i>Erithacus rubecula</i>)	1
10	European Serin (<i>Serinus serinus</i>)	1
11	Fieldfare (<i>Turdus pilaris</i>)	1
12	Greenfinch (<i>Carduelis chloris</i>)	1
13	Green Turaco (<i>Tauraco persa</i>)	1
14	Purple Grnadier (<i>Granatina ianthinogaster</i>)	2
15	Red-tailed Longhing Thrush (<i>Trochalopteron milnei</i>)	1
16	White-bellied Go-away Bird (<i>Criniferoides leucogaster</i>)	1
17	Wood Sandpiper (<i>Tringa glareola</i>)	1

total: 17 Species of birds 27 detections